

# THE ROLE OF FOURIER MODES IN EXTENSION THEOREMS OF HARTOGS-CHIRKA TYPE

DAVID E. BARRETT AND GAUTAM BHARALI

ABSTRACT. We generalize Chirka's theorem on the extension of functions holomorphic in a neighbourhood of  $\Gamma(F) \cup (\partial D \times D)$  – where  $D$  is the open unit disc and  $\Gamma(F)$  is the graph of a continuous  $D$ -valued function  $F$  – to the bidisc. We extend holomorphic functions by applying the Kontinuitätssatz to certain continuous families of analytic annuli, which is a procedure suited to configurations not covered by Chirka's theorem.

## 1. INTRODUCTION AND STATEMENT OF RESULTS

This article is motivated by the paper [3] by Chirka, in which the following theorem is proved (in what follows,  $D$  will represent the open unit disc in  $\mathbb{C}$  with centre at the origin, and given a function  $F$  defined in some region in  $\mathbb{C}$ ,  $\Gamma(F)$  will denote the graph of  $F$  over its domain) :

**Theorem (Chirka).** *Let  $F \in \mathcal{C}(\overline{D}; \mathbb{C})$  and assume that  $\sup_{\overline{D}} |F| < 1$ . Let  $\Omega$  be a connected neighbourhood of  $\Gamma(F) \cup (\partial D \times D)$  contained in  $\mathbb{C} \times D$ . If  $f \in \mathcal{O}(\Omega)$ , then  $f$  extends holomorphically to the bidisc  $D \times D$ .*

The requirement that  $\sup_{\overline{D}} |F| < 1$  is rather essential to the extension theorem stated above (in contrast, refer to [4] for a version by Chirka & Rosay, in which the condition  $\sup_{\overline{D}} |F| < 1$  is relaxed, but in which only the functions holomorphic in the union of a neighbourhood of  $\Gamma(F)$  with  $\{z \in \mathbb{C} : |z| > 1\} \times D$  – i.e. holomorphic in a *large* domain – extend holomorphically). A pertinent counterexample, when  $\sup_{\overline{D}} |F| > 1$ , to the sort of holomorphic extension described in Chirka's theorem – i.e. extension from *small* neighbourhoods of  $\Gamma(F) \cup (\partial D \times D)$  – is the case when  $\Gamma(F)$  is a Wermer disc. We will discuss this example in §4 below.

The strategy of Chirka – inspired by the methods in [7] – is to construct a continuously varying family of functions  $\{F_t\}_{t \in [0,1]} \subset \{G \in \mathcal{C}(\mathbb{C}) \mid \lim_{|z| \rightarrow \infty} G(z) = 0\}$  such that  $F_1 = \widetilde{F}$  and  $F_0 \equiv 0$ , and such that  $\Gamma(F_t)$  is complex-analytic in a neighbourhood of any  $(z, F_t(z)) \notin \Omega \cup (\{ |z| > 1\} \times D)$ . Here,  $\widetilde{F}$  is any smooth extension of the  $F$  provided by the theorem, that satisfies  $\widetilde{F}|_{|z| \geq 2} \equiv 0$ . Next, one extends  $f \in \mathcal{O}(\Omega)$  to  $\Omega \cup (\{ |z| > 1\} \times D)$  via Laurent decomposition. One can now show that the latter can be analytically continued, owing to the Kontinuitätssatz, via  $\{F_t\}_{t \in [0,1]}$  to a neighbourhood of the classical Hartogs configuration  $\Gamma(F_0) \cup (\partial D \times D)$ . The condition  $\sup_{\overline{D}} |F| < 1$  is crucial in ensuring that the extension of  $f \in \mathcal{O}(\Omega)$  by Laurent decomposition is single-valued. The strategy described fails in  $\mathbb{C}^n$ ,  $n > 2$ , and Chirka's theorem does not extend to higher dimensions as shown by Rosay's counterexample in [8].

---

2000 *Mathematics Subject Classification.* Primary 32D15.

*Key words and phrases.* Hartogs-Chirka type extension, holomorphic extension.

The work of the first author is supported by NSF Grant DMS-0072237.

Appeared in *Math. Z.* **249** (2005), 883-901.

The results in this paper are motivated by the two-fold aim of :

- a) Showing that functions holomorphic in *small* neighbourhoods of a Hartogs-Chirka type configuration  $\Gamma(F) \cup (\partial D \times D)$ , with  $\sup_{\overline{D}} |F| \gg 1$ , extend holomorphically to  $D \times D$  (in a manner that will be made precise in Theorem 1.1), given that  $F$  satisfies suitable restrictions.
- b) Extending Chirka's theorem to higher dimensions, and to a reasonably wide class of Hartogs-Chirka type configurations  $\Gamma(F) \cup (\partial D \times D^m)$ ,  $m \geq 2$  (in particular, to configurations in which  $F$  is not *merely* real-analytic or  $\mathcal{C}^\infty$ ).

Neither of the above seems to be achievable using Chirka's strategy. In this article, we discuss an alternative strategy for invoking the Kontinuitätssatz, and use it to demonstrate new Hartogs-Chirka type extension phenomena.

The first of the above aims is met by the following theorem. But we first present the following notation : if  $\Omega$  is a domain in  $\mathbb{C}^n$ , then  $(\widetilde{\Omega}, \pi^\Omega)$  will denote the envelope of holomorphy of  $\Omega$ .

**Theorem 1.1.** *Let  $F \in \mathcal{C}(\overline{D}; \mathbb{C})$  and assume that  $\sup_{\partial D} |F| < 1$ . Let  $A_j(r)$  represent the  $j^{\text{th}}$  Fourier coefficient of  $F(re^{i\cdot})$ ,  $r > 0$ ,  $j \in \mathbb{Z}$ . Assume that  $F$  satisfies the condition*

$$(1.1) \quad \sum_{n \in \mathbb{Z}} \frac{|A_n(r)|}{r^n} < 1 \quad \forall r \in (0, 1].$$

*Let  $\Omega_1$  be a neighbourhood of  $\Gamma(F) \cup (\partial D \times D)$  and let  $\Omega_2$  be any connected open set satisfying  $\partial D \times D \subset \overline{\Omega_2} \subset \Omega_1 \cap (\{|z| \geq 1\} \times D)$ . If  $f \in \mathcal{O}(\Omega_1)$ , then  $f|_{\Omega_2}$  has a holomorphic extension to  $D \times D$ .*

Note that when  $\sup_{\overline{D}} |F| > 1$ ,  $\pi^{\Omega_1}(\widetilde{\Omega_1}) \not\supseteq D \times D$  in general. For this reason, the usual arguments justifying that  $f$  has a single-valued extension to the bidisc fail. This is the reason behind the particular form of the conclusion of Theorem 1.1. Observe that while the condition (1.1) admits  $F$  such that the negative Fourier modes of  $F(re^{i\cdot})$  are large, it imposes a severe restriction on the sizes of the positive Fourier modes of  $F(re^{i\cdot})$  as  $r \rightarrow 0^+$ . One would like to investigate if such severe restrictions on the positive Fourier modes are necessary. This is a valid concern because if we assume that the function  $F$  has *only* positive Fourier modes, the condition (1.1) becomes unnecessary. The relevant theorem in this case is

**Theorem 1.2.** *Let  $F \in \mathcal{C}(\overline{D}; \mathbb{C})$  and assume that  $\sup_{\partial D} |F| < 1$ . Let  $A_j(r)$  represent the  $j^{\text{th}}$  Fourier coefficient of  $F(re^{i\cdot})$ ,  $r > 0$ ,  $j \in \mathbb{Z}$ . Assume that  $A_j \equiv 0 \quad \forall j < 0$ . Let  $\Omega_1$  be a neighbourhood of  $\Gamma(F) \cup (\partial D \times D)$  and let  $\Omega_2$  be any connected open set satisfying  $\partial D \times D \subset \overline{\Omega_2} \subset \Omega_1 \cap (\{|z| \geq 1\} \times D)$ . If  $f \in \mathcal{O}(\Omega_1)$ , then  $f|_{\Omega_2}$  has a holomorphic extension to  $D \times D$ .*

In a somewhat different direction, we may consider a continuous mapping  $F := (F_1, \dots, F_m) : \overline{D} \rightarrow D^m$ ,  $m \geq 2$ , and consider the Hartogs-Chirka type configuration  $\Gamma(F) \cup (\partial D \times D^m)$ . We know that, in general, Chirka's result is not true for such higher-dimensional configurations – see [8]. In contrast, it has been shown by Bharali [2] that Chirka's result does generalize to a certain class of Hartogs-Chirka type configurations. However, the class of real-analytic maps  $(F_1, \dots, F_m)$  studied in [2] is rather restrictive. We show in this paper that that if we impose a condition analogous to condition (1.1) above, we can demonstrate analytic continuation for a considerably less restrictive set of configurations. We make this precise in the following

**Theorem 1.3.** *Let  $F = (F_1, \dots, F_m) \in \mathcal{C}(\overline{D}; \mathbb{C}^m)$ . Assume that  $F(e^{i\theta}) \in D^m \forall \theta \in [0, 2\pi)$  and let  $A_{jk}(r)$  represent the  $k^{\text{th}}$  Fourier coefficient of  $F_j(re^{i\cdot})$ ,  $r > 0$ ,  $k \in \mathbb{Z}$ ,  $j = 1, \dots, m$ . Assume that each  $F_j$  satisfies the condition*

$$(1.2) \quad \sum_{n \in \mathbb{Z}} \frac{|A_{jn}(r)|}{r^n} < 1 \quad \forall r \in (0, 1].$$

*Let  $\Omega_1$  be a neighbourhood of  $\Gamma(F) \cup (\partial D \times D^m)$  and let  $\Omega_2$  be any connected open set satisfying  $\partial D \times D^m \subset \overline{\Omega_2} \subset \Omega_1 \cap (\{|z| \geq 1\} \times D^m)$ . If  $f \in \mathcal{O}(\Omega_1)$ , then  $f|_{\Omega_2}$  has a holomorphic extension to  $D \times D^m$ .*

We note that if, in Theorem 1.3,  $F$  were to satisfy the restriction  $F(\zeta) \in D^m \forall \zeta \in \overline{D}$ , then all functions  $f \in \mathcal{O}(\Omega)$  – where  $\Omega$  is a connected neighbourhood of  $\Gamma(F) \cup (\partial D \times D^m)$  contained in  $\mathbb{C} \times D^m$  – would extend to  $D^m$ , which is just Chirka’s extension phenomenon in a restricted, higher-dimensional setting.

The approach used in the first and the third theorem is to construct a continuous family of analytic annuli which are attached to  $\Gamma(H)$  – where  $H$  is an appropriately selected perturbation of  $F$  – along their inner boundaries, and to  $\partial D \times D^m$  (with  $m = 1$  in Theorem 1.1 and  $m \geq 2$  in Theorem 1.3) along their outer boundaries. Once this family is constructed, analytic continuation is achieved by invoking the Kontinuitätssatz. The proof of Theorem 1.2 uses a similar idea, but involves continuous families of analytic discs. These proofs may be found in §3. The technical construction of the aforementioned families of annuli/discs is carried out in the next section.

In the final section of this paper, we discuss a few examples. Firstly, we show that one can construct Hartogs-Chirka type configurations  $\Gamma(F) \cup (\partial D \times D)$  such that  $\sup_{\overline{D}} |F|$  is as large as we want and such that functions holomorphic in small neighbourhoods of this configuration extend. Next, we discuss a configuration involving Wermer’s disc (see Example 4.2 for a definition) – for which the extension phenomenon occurring in the previous example fails. And lastly, we show how Rosay’s counterexample to a higher-dimensional analogue of Chirka’s theorem fails to satisfy the hypotheses of Theorem 1.3.

## 2. PRELIMINARY LEMMAS

We need a few preliminary lemmas before we can prove our main theorems. In what follows,  $\text{Ann}(a; r, R)$  will denote the open annulus with centre at  $a \in \mathbb{C}$  and having inner and outer radii  $r$  and  $R$  respectively, while  $D(a; R)$  will denote the open disc of radius  $R$  with centre at  $a$ . The symbol  $\mathcal{C}^\infty(\overline{D}; \mathbb{C}^m)$ ,  $m = 1, 2, \dots$ , will denote the class of infinitely differentiable functions on the unit disc, all of whose derivatives extend to continuous functions on  $\overline{D}$ .

The reader will notice that in the following lemma the hypothesis  $G \in \mathcal{C}^\infty(\overline{D}; \mathbb{C})$  is much stronger than is required for the conclusion of Lemma 2.1. The only place where we use this hypothesis is in showing the existence of a certain limit towards the end of the proof. However, stating the strongest versions of Lemmas 2.1 and 2.3 – which are of relatively minimal utility in themselves – merely results in statements that are overly technical. For this reason, the  $G$  occurring in Lemmas 2.1-2.4 shall be assumed to be  $\mathcal{C}^\infty$ .

**Lemma 2.1.** *Let  $G(re^{i\theta}) = \sum_{n=-N}^N b_n(r)e^{in\theta}$  and assume that  $G \in \mathcal{C}^\infty(\overline{D}; \mathbb{C})$ . Assume further that*

$$(2.1) \quad \sum_{n=-N}^N \frac{|b_n(r)|}{r^n} < 1 \quad \forall r \in (0, 1].$$

*Then the holomorphic function*

$$\mathcal{A}_r(\zeta) = \sum_{n=-N}^N b_n(r) \left( \frac{\zeta}{r} \right)^n, \quad \zeta \in \text{Ann}(0; r, 1),$$

*which belongs to  $\mathcal{O}[\text{Ann}(0; r, 1)] \cap \mathcal{C}[\overline{\text{Ann}}(0; r, 1)]$ , satisfies  $|\mathcal{A}_r(e^{i\theta})| < 1$ . Fix  $\nu \in \mathbb{N}$  and let  $K \Subset \text{Ann}(0; 1/\nu, 1)$  be a compact subset. The function  $(0, 1/\nu] \times K \ni (r, \zeta) \mapsto \mathcal{A}_r(\zeta)$  extends to a continuous function on  $[0, 1/\nu] \times K$ .*

*Proof.* To prove the first part of this lemma, note that

$$(2.2) \quad |\mathcal{A}_r(e^{i\theta})| \leq \sum_{n=-N}^N |b_n(r)| \left| \frac{e^{i\theta}}{r} \right|^n = \sum_{n=-N}^N \frac{|b_n(r)|}{r^n} < 1.$$

We fix  $\nu \in \mathbb{N}$  and then fix a compact set  $K \Subset \text{Ann}(0; 1/\nu, 1)$ . It is obvious that  $(0, \nu] \times K \ni (r, \zeta) \mapsto b_{-n}(r)(r/\zeta)^n$  extends to a continuous function on  $[0, 1/\nu] \times K$ , which simply vanishes when  $r = 0$ , for each  $n = 1, 2, \dots, N$ . Now consider the function  $(r, \zeta) \mapsto b_n(r)(\zeta/r)^n$ ,  $n = 1, 2, \dots, N$ . Note that (2.1)  $\implies |b_n(r)| < r^n \forall n = 1, 2, \dots, N$ . This implies, since  $G$  is assumed to be smooth, that each of the latter functions extends continuously to a function  $\varphi_n \in \mathcal{C}([0, 1/\nu] \times K)$ , which is defined as

$$\varphi_n(r, \zeta) := \begin{cases} b_n(r)(\zeta/r)^n, & \text{if } (r, \zeta) \in (0, 1/\nu] \times K, \\ \frac{1}{n!} \left. \frac{d^n b_n}{dr^n} \right|_{r=0} \zeta^n, & \text{if } (r, \zeta) \in \{0\} \times K. \end{cases}$$

Since  $\mathcal{A}_r$  is a finite sum of the functions  $b_n(r)(r/\zeta)^n$ , the last two observations establish the second part of this lemma.  $\square$

**Lemma 2.2.** *Let  $G$  be as in Lemma 2.1, but assume additionally that  $\sup_{\partial D} |G| < 1$ . Let  $\Omega_1$  be a neighbourhood of  $\Gamma(G) \cup (\partial D \times D)$  such as that described in Theorem 1.1. Then*

- $\{\mathcal{A}_r\}_{r \in (0, 1)}$  is a continuous family in the sense that for a fixed  $\zeta_0 \in D \setminus \{0\}$ ,  $r \mapsto \mathcal{A}_r(\zeta_0)$  is continuous in the interval  $(0, |\zeta_0|)$ .
- $\lim_{r \rightarrow 0^+} \mathcal{A}_r(\zeta_0)$  exists for each  $\zeta_0 \in D \setminus \{0\}$ , and there exists a  $\psi \in \mathcal{O}(D)$  such that  $\psi(\zeta) = \lim_{r \rightarrow 0^+} \mathcal{A}_r(\zeta)$  on  $D \setminus \{0\}$ .
- Define

$$\mathfrak{K} := \Gamma(\psi) \cup \left[ \cup_{0 < r < 1} \{(\zeta, \mathcal{A}_r(\zeta)) \in \mathbb{C}^2 \mid r < |\zeta| < 1\} \right] \setminus \Omega_1.$$

$\mathfrak{K}$  is compact.

*Proof.* Part (a) and the first half of part (b) are obvious conclusions of Lemma 2.1. Thus, we may define

$$\psi(\zeta) := \lim_{|\zeta| > r \rightarrow 0^+} \mathcal{A}_r(\zeta) \quad \forall \zeta \in D \setminus \{0\}.$$

Fix a  $\nu \in \mathbb{N}$ . Lemma 2.1(b) tells us that

$$(\mathcal{A}_r|_{\text{Ann}(0; 1/\nu, 1)})(\zeta) \longrightarrow \psi(\zeta) \text{ uniformly on each compact } K \Subset \text{Ann}(0; 1/\nu, 1) \text{ as } r \searrow 0.$$

We conclude from this statement that

$$(2.3) \quad \psi|_{\text{Ann}(0;1/\nu,1)} \in \mathcal{O}[\text{Ann}(0;1/\nu,1)] \quad \forall \nu = 2, 3, 4, \dots$$

Before proceeding any further, we comment that the functions  $\mathcal{A}_r$  are so constructed that  $\Gamma(\mathcal{A}_r)$ ,  $0 < r < 1$ , are analytic annuli that are attached to  $\Gamma(G)$  along their inner boundaries and – in view of the inequality (2.2) – to  $\partial D \times D$  along their outer boundaries. Therefore,

$$(2.4) \quad |\mathcal{A}_r(\zeta)| \leq \max \left\{ \sup_{|\xi|=r} |\mathcal{A}_r(\xi)|, 1 \right\} \leq \max \left\{ \sup_{\overline{D}} |G|, 1 \right\} \\ \forall \zeta \in \text{Ann}(0;r,1) \text{ and for each } r \in (0,1).$$

By (2.3),  $\psi$  is already holomorphic on  $D \setminus \{0\}$ . The bounds above imply, since  $\psi(\zeta)$  is the limit of the  $\mathcal{A}_r(\zeta)$ 's, provided  $\zeta \neq 0$ , that  $|\psi(\zeta)| \leq \sup_{\xi \neq 0} |G(\xi)|$  in a punctured neighbourhood of the origin. Thus,  $\psi$  extends to a holomorphic function on  $D$ . This establishes (b).

Notice that by the estimates (2.4) and part (b) of this lemma,  $\mathfrak{K}$  is a bounded set. Therefore, it suffices to show that  $\mathfrak{K}$  is closed. Now consider a point  $(z, w) \notin \Omega_1$  with the property that there exist sequences  $\{r(\nu)\}_{\nu \in \mathbb{N}} \subset (0, 1)$  and  $\{\zeta_\nu\}_{\nu \in \mathbb{N}} \subset D$  such that  $r(\nu) \rightarrow 0$  as  $\nu \rightarrow \infty$  and  $(\zeta_\nu, \mathcal{A}_{r(\nu)}(\zeta_\nu)) \rightarrow (z, w)$  as  $\nu \rightarrow \infty$ . It is easy to see that to prove (c), it suffices to show that all such points  $(z, w) \in \mathfrak{K}$ . Notice that, by construction, there is a  $\delta(\Omega_1) > 0$  depending only on  $\Omega_1$  such that  $(\zeta, \mathcal{A}_r(\zeta)) \in \Omega_1 \forall r, |\zeta| < \delta(\Omega_1)$ . Thus, as  $(z, w) \notin \Omega_1$ ,  $z \neq 0$ . Now, given that the  $\mathcal{A}_r$ 's converge uniformly on compact subsets lying away from 0, there exists  $\kappa_1 \in \mathbb{N}$  such that

$$|\mathcal{A}_{r(\nu)}(\zeta) - \psi(\zeta)| < \varepsilon/2 \quad \forall \nu \geq \kappa_1, \forall \zeta \in \overline{D}(z; |z|/2).$$

Let  $\kappa_2 \in \mathbb{N}$  be such that

$$\zeta_\nu \in D(z; |z|/2) \quad \text{and} \quad |w - \mathcal{A}_{r(\nu)}(\zeta_\nu)| < \varepsilon/2 \quad \forall \nu \geq \kappa_2.$$

The above inequalities imply that

$$|w - \psi(\zeta_\nu)| \leq |w - \mathcal{A}_{r(\nu)}(\zeta_\nu)| + |\mathcal{A}_{r(\nu)}(\zeta_\nu) - \psi(\zeta_\nu)| < \varepsilon \quad \forall \nu \geq \max(\kappa_1, \kappa_2).$$

This tells us that  $w = \lim_{\nu \rightarrow \infty} \psi(\zeta_\nu)$ , whence  $(z, w) \in \Gamma(\psi) \setminus \Omega_1$ . This establishes (c), and concludes our proof.  $\square$

**Lemma 2.3.** *Let  $G(re^{i\theta}) = \sum_{n=0}^N b_n(r)e^{in\theta}$  – i.e. we assume that  $G(re^{i\cdot})$  has no negative Fourier modes. Assume further that  $G \in \mathcal{C}^\infty(\overline{D}; \mathbb{C})$ . Then the holomorphic function*

$$\mathcal{D}_r(\zeta) = \sum_{n=0}^N b_n(r) \left( \frac{\zeta}{r} \right)^n, \quad \zeta \in D(0; r),$$

*which belongs to  $\mathcal{O}[D(0; r)] \cap \mathcal{C}[\overline{D}(0; r)]$ , satisfies  $\mathcal{D}_r(re^{i\theta}) = G(re^{i\theta}) \forall \theta \in [0, 2\pi)$ . Fix  $\nu \in \mathbb{N}$  and let  $K \Subset D(0; 1 - 1/\nu)$  be a compact subset. The function  $(r, \zeta) \mapsto \mathcal{D}_r(\zeta)$  is a continuous function on  $[1 - 1/\nu, 1] \times K$ .*

The above lemma is a triviality; we merely state it as an element that will be needed in the proof of our next result.

**Lemma 2.4.** *Let  $G$  be as in Lemma 2.3, but assume additionally that  $\sup_{\partial D} |G| < 1$ . Let  $\Omega_1$  be a neighbourhood of  $\Gamma(G) \cup (\partial D \times D)$  such as that described in Theorem 1.2. Then*

- a)  $\{\mathcal{D}_r\}_{r \in (0,1)}$  is a continuous family in the sense that for a fixed  $\zeta_0 \in D$ ,  $r \mapsto \mathcal{D}_r(\zeta_0)$  is continuous in the interval  $(|\zeta_0|, 1)$ .
- b)  $\lim_{r \rightarrow 1^-} \mathcal{D}_r(\zeta)$  exists for each  $\zeta \in D$ , and this limit defines a holomorphic function  $\psi \in \mathcal{O}(D)$ .
- c) Define

$$\mathfrak{K} := \Gamma(\psi) \cup \left[ \cup_{0 < r < 1} \{(\zeta, \mathcal{D}_r(\zeta)) \in \mathbb{C}^2 \mid |\zeta| < r\} \right] \setminus \Omega_1.$$

$\mathfrak{K}$  is compact.

*Proof.* Part (a) and the first half of part (b) are direct consequences of Lemma 2.3. The inference that

$$\psi(\zeta) := \lim_{|\zeta| < r \rightarrow 1^-} \mathcal{D}_r(\zeta) \quad \forall \zeta \in D$$

is holomorphic follows from Lemma 2.3. The uniform-convergence argument is exactly analogous to the argument used in proving Lemma 2.2. We therefore omit the details. We remark that

$$\psi(\zeta) = \sum_{n=0}^N b_n(1)\zeta^n.$$

The functions  $\mathcal{D}_r$  are so constructed that  $\Gamma(\mathcal{D}_r)$ ,  $0 < r < 1$ , are analytic discs that are attached to  $\Gamma(G)$  along their boundaries. Therefore,

$$|\mathcal{D}_r(\zeta)| \leq \sup_{|\xi|=r} |\mathcal{D}_r(\xi)| \leq \sup_{\overline{D}} |G| \quad \forall \zeta \in D(0; r) \text{ and for each } r \geq 1 - 1/\nu.$$

Thus,  $\mathfrak{K}$  is a bounded set, and we argue that  $\mathfrak{K}$  is closed exactly as we did in Lemma 2.2(c).  $\square$

The following lemma is key to the proofs of Theorems 1.1-1.3. Before proving it, we explicitly state the following simple

**Fact :** *Due to the continuity of the functions  $F$  and  $G$  occurring in the statements of the various theorems and lemmas above, the associated Fourier coefficients  $A_n(r)$  and  $b_n(r)$  satisfy  $A_n(0) = b_n(0) = 0 \forall n \neq 0$ .*

This fact is used implicitly at several places in the next lemma.

**Lemma 2.5.** *Let  $F \in \mathcal{C}(\overline{D}; \mathbb{C})$  and let  $A_j(r)$  represent the  $j^{\text{th}}$  Fourier coefficient of  $F(re^{i\cdot})$ ,  $r > 0$ ,  $j \in \mathbb{Z}$ . Assume that :*

- 1)  $\sup_{\partial D} |F| < 1$ , and
- 2)  $F$  satisfies the condition

$$(2.5) \quad \sum_{n \in \mathbb{Z}} \frac{|A_n(r)|}{r^n} < 1 \quad \forall r \in (0, 1].$$

Given  $\varepsilon > 0$  there exists a function  $G \in \mathcal{C}^\infty(\overline{D}; \mathbb{C})$  of the form

$$G(re^{i\theta}) = \sum_{n=-N}^N B_n(r)e^{in\theta},$$

where  $N$  is some large positive integer and  $B_n \in \mathcal{C}^\infty([0, 1]; \mathbb{C})$ , such that

- $|F(\zeta) - G(\zeta)| < \varepsilon \forall \zeta \in \overline{D}$ ,
- $G$  has the property (1) and satisfies the analogue of (2) above (with  $B_n(r)$  replacing  $A_n(r)$  in (2.5) above).

Furthermore, if property (2) is replaced by

2\*)  $F$  has no negative Fourier modes,

then  $G$  can be constructed so that it has property (1) and  $B_{-j} \equiv 0$  for  $j = 1, 2, \dots, N$ .

*Proof.* Define

$$S_m(\theta, r) := \sum_{j=-m}^m A_j(r) e^{ij\theta}$$

$$\sigma_n(\theta, r) := \frac{S_0(r, \theta) + \dots + S_n(\theta, r)}{n+1}$$

Let us first assume that  $F$  has properties (1) and (2). Let  $\eta > 0$  be so small that

$$(2.6) \quad \eta < 1 - \sup_{\partial D} |F|,$$

$$\eta + \sum_{n \in \mathbb{Z}} |A_n(r)|/r^n < 1 \quad \forall r \in (0, 1],$$

and define  $\delta := \min(\varepsilon, \eta)$ . There exists a natural number  $N > 0$  such that

$$(2.7) \quad |F(re^{i\theta}) - \sigma_N(\theta, r)| < \delta/2 \quad \forall (\theta, r) \in [0, 2\pi) \times [0, 1].$$

This above is a consequence of Fejér's theorem. For a fixed  $r \in [0, 1]$ , (2.7) is precisely the statement of Fejér's theorem applied to the periodic function  $F(re^{i\cdot})$ . However, on examining the proof of Fejér's theorem, one sees that owing to the equicontinuity of the family  $\{F(re^{i\cdot})\}_{r \in [0, 1]} \subset \mathcal{C}(\mathbb{T})$ , the choice of  $N$  in (2.7) is uniform in  $r \in [0, 1]$ .

One sees immediately that if one writes

$$\sigma_N(\theta, r) = \sum_{j=-N}^N a_j(r) e^{ij\theta},$$

then  $|a_j(r)| \leq |A_j(r)| \forall r \in [0, 1]$ . For each  $j = 1, 2, \dots, N$ , we pick a function  $B_{-j}(r)$  which satisfies the following conditions :

- i)  $B_{-j} \in \mathcal{C}^\infty([0, 1]; \mathbb{C})$ ,
- ii)  $B_{-j}$  vanishes to infinite order at  $r = 0$ , and
- iii)  $|a_{-j}(r) - B_{-j}(r)| \leq \delta/2(2N+1) \forall r \in [0, 1]$ ,

provided  $a_{-j} \not\equiv 0$ ,  $j = 1, 2, \dots, N$ . If  $a_{-j} \equiv 0$ , we just choose  $B_{-j} \equiv 0$ . Note that by our condition on the Fourier coefficients  $\{A_n(r)\}_{n \in \mathbb{Z}}$ ,  $|a_j(r)| \leq |A_j(r)| \leq r^j \forall j = 1, 2, \dots, N$ ,  $r \in [0, 1]$ . Let  $0 < R_0 < 1$  be a small number such that

$$R_0 \leq \left\{ \frac{\delta}{4(2N+1)} \right\}^{1/j} \quad \forall j = 1, 2, \dots, N.$$

For each  $j = 1, 2, \dots, N$  we define a function  $B_j(r)$  as follows :

$$B_j(r) := \begin{cases} \alpha_j(r) r^j, & \text{if } r \leq R_0, \\ \beta_j(r), & \text{if } r \geq R_0, \end{cases}$$

such that

- i\*)  $B_j \in \mathcal{C}^\infty([0, 1]; \mathbb{C})$ ,
- ii\*)  $\alpha_j$  vanishes to infinite order at  $r = 0$ ,

*iii\**)  $\alpha_j$  satisfies

$$|\alpha_j(r)| \leq \sup_{s \leq 1} \frac{|a_j(s)|}{s^j} \quad \forall s \in [0, R_0],$$

*iv\**)  $\beta_j$  satisfies

$$|\beta_j(r) - a_j(r)| \leq \frac{R_0^j \delta}{2(2N+1)} \quad \forall r \in [R_0, 1].$$

Finally, define  $B_0(r)$  to be any  $\mathcal{C}^\infty$  function such that  $|B_0(r) - a_0(r)| < \delta/2(2N+1) \forall r \in [0, 1]$  and such that  $B_0 - B_0(0)$  vanishes to high order at  $r = 0$ . Now write

$$G(re^{i\theta}) = \sum_{j=-N}^N B_j(r) e^{ij\theta}.$$

We now make some estimates. We first consider the  $B_{-j}(r)$ 's,  $j = 1, 2, \dots, N$ . Note that the following statements continue to be true trivially if  $B_{-j} \equiv 0$  for any  $j = 1, 2, \dots, N$ .

$$(2.8) \quad \sum_{j=1}^N |B_{-j}(r) - a_{-j}(r)| \leq \sum_{j=1}^N \frac{\delta}{2(2N+1)} \leq \frac{\delta N}{2(2N+1)},$$

$$(2.9) \quad \begin{aligned} \sum_{j=1}^N |B_{-j}(r)| r^j &\leq \sum_{j=1}^N r^j \left\{ |a_{-j}(r)| + \frac{\delta}{2(2N+1)} \right\} \\ &\leq \sum_{j=1}^N |a_{-j}(r)| r^j + \frac{\delta N}{2(2N+1)} \quad \forall r \in [0, 1]. \end{aligned}$$

Next, we consider the  $B_j(r)$ 's,  $j = 1, 2, \dots, N$ . First, we let  $0 \leq r \leq R_0$ . We use item (*iii\**) in the definition of  $B_j(r)$  above to get :

$$(2.10) \quad \begin{aligned} \sum_{j=1}^N |B_j(r) - a_j(r)| &\leq \sum_{j=1}^N r^j \left| \alpha_j(r) - \frac{|a_j(r)|}{r^j} \right| \\ &\leq \sum_{j=1}^N 2R_0^j \sup_{s \leq 1} \frac{|a_j(s)|}{s^j} \\ &\leq \sum_{j=1}^N 2 \left\{ \frac{\delta}{4(2N+1)} \right\} \sup_{s \leq 1} \frac{|A_j(s)|}{s^j} \\ &\leq \frac{\delta N}{2(2N+1)}, \end{aligned}$$

$$(2.11) \quad \sum_{j=1}^N \frac{|B_j(r)|}{r^j} \leq \sum_{j=1}^N \sup_{s \leq 1} \frac{|a_j(s)|}{s^j} \quad \forall r \in [0, R_0].$$



And when we consider  $R_0 \leq r \leq 1$ , we use item  $(iv^*)$  in the definition of  $B_j(r)$  to get :

$$(2.12) \quad \sum_{j=1}^N |B_j(r) - a_j(r)| \leq \sum_{j=1}^N \frac{R_0^j \delta}{2(2N+1)} \leq \frac{\delta N}{2(2N+1)},$$

$$(2.13) \quad \begin{aligned} \sum_{j=1}^N \frac{|B_j(r)|}{r^j} &\leq \sum_{j=1}^N \left\{ \frac{|a_j(r)|}{r^j} + \frac{R_0^j \delta}{2(2N+1)R_0^j} \right\} \\ &\leq \sum_{j=1}^N \frac{|a_j(r)|}{r^j} + \frac{\delta N}{2(2N+1)} \quad \forall r \in [R_0, 1]. \end{aligned}$$

Observe that from (2.9) and (2.11), we have

$$(2.14) \quad \begin{aligned} \sum_{j=-N}^N \frac{|B_j(r)|}{r^j} &\leq \sum_{j=1}^N |a_{-j}(r)| r^j + \frac{\delta N}{2(2N+1)} + |a_0(r)| + \frac{\delta}{2(2N+1)} + \sum_{j=1}^N \sup_{s \leq 1} \frac{|a_j(s)|}{s^j} \\ &\leq \sum_{j=-N}^N \sup_{s \leq 1} \frac{|a_j(s)|}{s^j} + \frac{\delta(N+1)}{2(2N+1)} \\ &\leq \sum_{j=-N}^N \sup_{s \leq 1} \frac{|A_j(s)|}{s^j} + \frac{\eta}{2} < 1 \quad \forall r \in [0, R_0]. \end{aligned}$$

The last inequality follows from the definition of  $\eta$  and the fact that  $|a_j(r)| \leq |A_j(r)| \forall r \in [0, 1]$ . Next, applying (2.9) and (2.13) we get

$$(2.15) \quad \begin{aligned} \sum_{j=-N}^N \frac{|B_j(r)|}{r^j} &\leq \sum_{j \neq 0} \frac{|a_j(r)|}{r^j} + \frac{2\delta N}{2(2N+1)} + |a_0(r)| + \frac{\delta}{2(2N+1)} \\ &\leq \sum_{j=-N}^N \sup_{s \leq 1} \frac{|a_j(s)|}{s^j} + \frac{\delta}{2} \\ &\leq \sum_{j=-N}^N \sup_{s \leq 1} \frac{|A_j(s)|}{s^j} + \frac{\eta}{2} < 1 \quad \forall r \in [R_0, 1]. \end{aligned}$$

From the inequalities (2.6), (2.14) and (2.15), we get

$$\sum_{j=-N}^N \frac{|B_j(r)|}{r^j} < 1 \quad \forall r \in [0, 1],$$

which is to say that  $G$  satisfies the analogue of (2), with  $B_n(r)$  replacing  $A_n(r)$  in the expression (2.5).

We now exploit the estimates (2.8), (2.10) and (2.12), to get

$$(2.16) \quad \begin{aligned} |F(re^{i\theta}) - G(re^{i\theta})| &\leq |F(re^{i\theta}) - \sigma_N(re^{i\theta})| + \sum_{j=-N}^N |a_j(r) - B_j(r)| \\ &< \frac{\delta}{2} + 2 \cdot \frac{\delta N}{2(2N+1)} + \frac{\delta}{2(2N+1)} = \delta. \end{aligned}$$

Given the way in which  $\delta$  is defined, we see that  $G$  has the property (1), and  $|G(\zeta) - F(\zeta)| < \varepsilon \forall \zeta \in \overline{D}$ .  $G$  is, of course, smooth by construction.

Note further that in the above construction, if  $F$  has no negative Fourier modes, neither does  $G$ . So, if  $F$  had the property (2\*) instead of property (2), in addition to choosing  $B_{-j} \equiv 0$  we would use the same rule by which we selected  $B_0(r)$  in the above argument to define  $B_j(r)$ ,  $j = 1, 2, \dots, N$ . It is easy to verify that this modified construction would yield the second part of this lemma.  $\square$

### 3. PROOFS OF THE THEOREMS

**3.1. The proof of Theorem 1.1.** Let  $\varepsilon > 0$  be so small that  $F(D) + \eta \subset \Omega_1 \forall \eta \in \mathbb{C}$  such that  $|\eta| < 2\varepsilon$ . By Lemma 2.5, there exists a function  $H \in \mathcal{C}^\infty(\overline{D}; \mathbb{C})$  with  $H(re^{i\theta}) = \sum_{n=-N}^N B_n(r)e^{in\theta}$  such that

$$|H(\zeta) - F(\zeta)| < \varepsilon \quad \forall \zeta \in \overline{D}, \quad \sup_{\partial D} |H| < 1,$$

$$\sum_{n=-N}^N \frac{|B_n(r)|}{r^n} < 1 \quad \forall r \in (0, 1].$$

Let  $\delta > 0$  be so small that

- $\sup_{\partial D} |H| + \delta < 1$ ;
- $\delta + \sum_{n=-N}^N |B_n(r)|/r^n < 1 \quad \forall r \in (0, 1]$ ; and
- $H(D) + \eta \subset \Omega_1 \quad \forall \eta \in \mathbb{C}$  such that  $|\eta| < \delta$ .

Define, for each  $\eta$  such that  $|\eta| < \delta$

$$H^{(\eta)}(\zeta) := H(\zeta) + \eta,$$

$$\mathcal{A}_r^{(\eta)}(\zeta) := \sum_{n \neq 0} B_n(r) \left(\frac{\zeta}{r}\right)^n + (B_0(r) + \eta), \quad \zeta \in \text{Ann}(0; r, 1),$$

We apply Lemma 2.2 to  $\{\mathcal{A}_r^{(\eta)}\}_{r \in (0, 1)}$  for each  $\eta$ , by taking

$$b_n(r) = B_n(r) \quad \forall n \neq 0, \quad b_0(r) = B_0(r) + \eta$$

in that lemma. We conclude that there is a function  $\psi \in \mathcal{O}(D)$  such that

- 1) For any fixed  $\zeta_0 \in D \setminus \{0\}$ ,  $r \mapsto \mathcal{A}_r^{(\eta)}(\zeta_0)$  is continuous in  $(0, |\zeta_0|)$  for  $|\eta| < \delta$ .
- 2) For  $\zeta \in D \setminus \{0\}$ ,  $\lim_{r \rightarrow 0^+} \mathcal{A}_r^{(\eta)}(\zeta) = \psi(\zeta) + \eta$  for  $|\eta| < \delta$ .
- 3) For each  $\eta : |\eta| < \delta$ ,  $\mathfrak{K}^{(\eta)}$  is compact, where we define

$$\mathfrak{K}^{(\eta)} := \Gamma(\psi + \eta) \cup \left[ \cup_{0 < r < 1} \{(\zeta, \mathcal{A}_r^{(\eta)}(\zeta)) \in \mathbb{C}^2 \mid r < |\zeta| < 1\} \right] \setminus \Omega_1.$$

In other words, for each fixed  $\eta$ , the family  $\{\Gamma(\mathcal{A}_r^{(\eta)})\}_{r \in (0, 1)}$  is a continuous family of analytic annuli attached to  $\Gamma(H^{(\eta)}) \cup (\partial D \times D)$ , which accumulate onto  $\Gamma(\psi + \eta)$  as  $r \rightarrow 0^+$ . The analytic annuli  $\Gamma(\mathcal{A}_r^{(\eta)})$  with  $r \approx 1$  are contained in  $\Omega_1$ . We can therefore apply the Kontinuitätssatz to conclude that

$$(3.1) \quad U(\Delta) := \{(z, w) \in D \times \mathbb{C} : (w - \psi(z)) \in \Delta\} = \bigcup_{\eta \in \Delta} \Gamma(\psi + \eta) \subset \pi^{\Omega_1}(\widetilde{\Omega}_1),$$

where  $\Delta$  is any disc contained in  $D(0; \delta)$

There is a canonical holomorphic imbedding of  $\Omega_1$  into  $\widetilde{\Omega}_1$ . We denote this imbedding by  $j : \Omega_1 \hookrightarrow \widetilde{\Omega}_1$ . Corresponding to each  $f \in \mathcal{O}(\Omega_1)$  there is a holomorphic function on  $\widetilde{\Omega}_1$ , which we

shall denote by  $\mathcal{E}(f)$ , such that  $\mathcal{E}(f) \circ j = f$ . It is now a standard argument – see, for instance [5] or [3] – to show that there exist holomorphic mappings

$$\widetilde{\mathcal{A}}_r(\cdot; \eta) : \text{Ann}(0; r, 1) \rightarrow \widetilde{\Omega}_1 \quad \forall r \in (0, 1), \quad \text{and} \quad \widetilde{\psi}(\cdot; \eta) : D \rightarrow \widetilde{\Omega}_1$$

such that

a) For each  $\eta$  with  $|\eta| < \delta$  :

$$\pi^{\Omega_1} \circ \widetilde{\mathcal{A}}_r(\zeta; \eta) = (\zeta, \mathcal{A}_r^{(\eta)}(\zeta)) \quad \forall \zeta \in \text{Ann}(0; r, 1) \text{ when } r \in (0, 1), \text{ and}$$

$$\pi^{\Omega_1} \circ \widetilde{\psi}(\zeta; \eta) = (\zeta, \psi(\zeta) + \eta) \quad \forall \zeta \in D.$$

b)  $j(\zeta, \mathcal{A}_r^{(\eta)}(\zeta)) = \widetilde{\mathcal{A}}_r(\zeta; \eta)$  wherever the left-hand side is defined, and  $\forall r \in (0, 1)$ .

c)  $j(\zeta, \psi(\zeta) + \eta) = \widetilde{\psi}(\zeta; \eta)$  wherever the left-hand side is defined.

Notice that – in view of item (a) – for each fixed  $\zeta \in D$ ,  $\eta \mapsto \widetilde{\psi}(\zeta; \eta)$  is holomorphic.

Let  $V$  be the connected component of  $(\pi^{\Omega_1})^{-1}(U(|\eta| < \delta))$  containing  $\mathcal{C}_0 := \text{image}(\widetilde{\psi}(\cdot; 0))$ . For each point  $q \in \mathcal{C}_0$  there is a neighbourhood  $W(q) \Subset V$  of  $q$  such that  $\pi^{\Omega_1}|_{W(q)} : W(q) \rightarrow \mathbb{C}^2$  is a biholomorphism. Let  $\Delta_*$  be a disc centered at the origin that is so small that

$$\text{image}(\widetilde{\psi}(\cdot; \eta)) \subset \bigcup_{q \in \mathcal{C}_0} W(q) \quad \forall \eta \in \Delta_*.$$

We define  $\Omega^* := U(\Delta_*) \cup \omega_2$ ,  $\omega_2$  being a connected open set satisfying

- $\overline{\Omega_2} \subset \omega_2 \subset \Omega_1$ ; and
- $\omega_2 \cap U(\Delta_*)$  is connected,

where  $\Omega_2$  is as described in Theorem 1.1, and  $U(\Delta_*)$  is as defined by (3.1). Our goal is to map  $\Omega^*$  into  $\widetilde{\Omega}_1$  in such a way that this mapping extends  $j$ . This mapping will allow us – given any  $f \in \mathcal{O}(\Omega_1)$  – to extend  $f|_{\omega_2}$  to  $\Omega^*$ . But  $\Omega^*$  is a neighbourhood of a classical Hartogs configuration, whence  $f|_{\Omega_2}$  would extend to the bidisc. To this end, we define

$$\widetilde{j}(z, w) := \begin{cases} \widetilde{\psi}(z; w - \psi(z)), & \text{if } (z, w) \in U(\Delta_*), \\ j(z, w), & \text{if } (z, w) \in \omega_2. \end{cases}$$

Note that if  $(z, w) \in U(\Delta_*) \cap \omega_2$ , then, in view of item (c) above

$$(3.2) \quad \widetilde{\psi}(z; w - \psi(z)) = j(z, \psi(z) + \{w - \psi(z)\}) = j(z, w) \quad \forall (z, w) \in U(\Delta_*) \cap \omega_2.$$

Thus,  $\widetilde{j}$  is well-defined, and extends  $j$ . From our foregoing remarks,  $\widetilde{j}$  is holomorphic. Given any  $f \in \mathcal{O}(\Omega_1)$ , we define  $\widetilde{f} \in \mathcal{O}(\Omega_*)$  by  $\widetilde{f} := \mathcal{E}(f) \circ \widetilde{j}$ . In view of (3.2),  $\widetilde{f}|_{\omega_2} \equiv f|_{\omega_2}$ . Notice that  $\Omega_*$  is a neighbourhood of  $\Gamma(\psi) \cup (\partial D \times D)$ , which is the classical Hartogs configuration. Thus  $\widetilde{f}$  has a holomorphic extension to  $D \times D$ , whence  $f|_{\Omega_2}$  has a holomorphic extension to  $D \times D$ . This concludes our proof.  $\square$

**3.2. The proof of Theorem 1.2.** Since the proof of this theorem is similar to that of Theorem 1.1, we shall be brief. Let  $\varepsilon > 0$  be so small that  $F(D) + \eta \subset \Omega_1 \quad \forall \eta \in \mathbb{C}$  such that  $|\eta| < 2\varepsilon$ . By Lemma 2.5, there exists a function  $H \in \mathcal{C}^\infty(\overline{D}; \mathbb{C})$  with  $H(re^{i\theta}) = \sum_{n=0}^N B_n(r)e^{in\theta}$  such that

$$|H(\zeta) - F(\zeta)| < \varepsilon \quad \forall \zeta \in \overline{D}, \quad \sup_{\partial D} |H| < 1.$$

Recall that the  $H$  that has the two properties above can be so chosen that it has no negative Fourier modes. Let  $\delta > 0$  be so small that

- $\sup_{\partial D} |H| + \delta < 1$ ; and
- $H(D) + \eta \subset \Omega_1 \ \forall \eta \in \mathbb{C}$  such that  $|\eta| < \delta$ .

For each  $\eta$  such that  $|\eta| < \delta$ , we define

$$\begin{aligned} H^{(\eta)}(\zeta) &:= H(\zeta) + \eta, \\ \mathcal{D}_r^{(\eta)}(\zeta) &:= \sum_{n=1}^N B_n(r) \left(\frac{\zeta}{r}\right)^n + (B_0(r) + \eta), \quad \zeta \in D(0; r), \end{aligned}$$

We apply Lemma 2.4 to  $\{\mathcal{D}_r^{(\eta)}\}_{r \in (0;1)}$  for each  $\eta$ , by taking

$$b_n(r) = B_n(r) \quad \forall n = 1, 2, \dots, N, \quad b_0(r) = B_0(r) + \eta$$

in that lemma. For each fixed  $\eta$ , the family  $\{\Gamma(\mathcal{D}_r^{(\eta)})\}_{r \in (0,1)}$  is a continuous family of analytic discs which are attached to  $\Gamma(H^{(\eta)})$  along their boundaries, and which accumulate onto a holomorphic graph  $\Gamma(\psi + \eta)$  as  $r \rightarrow 1^-$ . Furthermore, for  $|\eta| < \delta$ , each

$$\mathfrak{K}^{(\eta)} := \Gamma(\psi + \eta) \cup \left[ \cup_{0 < r < 1} \{(\zeta, \mathcal{D}_r^{(\eta)}(\zeta)) \in \mathbb{C}^2 \mid |\zeta| < r\} \right] \setminus \Omega_1$$

is compact. We may therefore apply the *Kontinuitätssatz* to conclude that

$$(3.3) \quad U(\Delta) := \{(z, w) \in D \times \mathbb{C} : (w - \psi(z)) \in \Delta\} = \bigcup_{\eta \in \Delta} \Gamma(\psi + \eta) \subset \pi^{\Omega_1}(\widetilde{\Omega}_1),$$

where  $\Delta$  is any disc contained in  $D(0; \delta)$

Let  $j : \Omega_1 \hookrightarrow \widetilde{\Omega}_1$  be the canonical holomorphic imbedding of  $\Omega_1$  into  $\widetilde{\Omega}_1$ . As before, there exist holomorphic mappings

$$\widetilde{\mathcal{D}}_r(\cdot; \eta) : D(0; r) \rightarrow \widetilde{\Omega}_1 \quad \forall r \in (0, 1), \quad \text{and} \quad \widetilde{\psi}(\cdot; \eta) : D \rightarrow \widetilde{\Omega}_1$$

such that

- a) For each  $\eta$  with  $|\eta| < \delta$  :

$$\begin{aligned} \pi^{\Omega_1} \circ \widetilde{\mathcal{D}}_r(\zeta; \eta) &= (\zeta, \mathcal{D}_r^{(\eta)}(\zeta)) \quad \forall \zeta \in D(0; r) \text{ when } r \in (0, 1), \text{ and} \\ \pi^{\Omega_1} \circ \widetilde{\psi}(\zeta; \eta) &= (\zeta, \psi(\zeta) + \eta) \quad \forall \zeta \in D. \end{aligned}$$

- b)  $j(\zeta, \mathcal{D}_r^{(\eta)}(\zeta)) = \widetilde{\mathcal{D}}_r(\zeta; \eta)$  wherever the left-hand side is defined, and  $\forall r \in (0, 1)$ .

- c)  $j(\zeta, \psi(\zeta) + \eta) = \widetilde{\psi}(\zeta; \eta)$  wherever the left-hand side is defined.

Arguing exactly as in the proof of Theorem 1.1, we can find a disc  $\Delta_* \subset D(0; \delta)$ , centered at the origin, and an appropriate open set  $\omega_2$  satisfying  $\overline{\Omega}_2 \subset \omega_2 \subset \Omega_1$ , such that if we define

$$\widetilde{j}(z, w) := \begin{cases} \widetilde{\psi}(z; w - \psi(z)), & \text{if } (z, w) \in U(\Delta_*), \\ j(z, w), & \text{if } (z, w) \in \omega_2, \end{cases}$$

(where  $U(\Delta_*)$  is as defined by (3.3) above), then  $\widetilde{j}$  is holomorphic and well defined. Holomorphicity follows from (a) above, while (c) implies that

$$(3.4) \quad \widetilde{\psi}(z; w - \psi(z)) = j(z, \psi(z) + \{w - \psi(z)\}) = j(z, w) \quad \forall (z, w) \in U(\Delta_*) \cap \omega_2.$$

We define  $\Omega^* := U(\Delta_*) \cup \omega_2$ . As before, given any  $f \in \mathcal{O}(\Omega_1)$ , we define  $\tilde{f} \in \mathcal{O}(\Omega_*)$  by  $\tilde{f} := \mathcal{E}(f) \circ \tilde{j}$ . In view of (3.4),  $\tilde{f}|_{\omega_2} \equiv f|_{\omega_2}$ . Since  $\Omega_*$  is a neighbourhood of  $\Gamma(\psi) \cup (\partial D \times D)$ , i.e. is a classical Hartogs configuration,  $\tilde{f}$  has a holomorphic extension to  $D \times D$ , whence  $f|_{\Omega_2}$  has a holomorphic extension to  $D \times D$ . This concludes our proof.  $\square$

**3.3. The proof of Theorem 1.3.** The proof of Theorem 1.3 proceeds along the same lines as the proof of the first theorem. The essential difference is that we find an  $\varepsilon > 0$  such that  $F(D) + \eta \subset \Omega \forall \eta \in D(0; 2\varepsilon)^m$ , and then apply Lemma 2.5 to the pairs  $(F_1, \varepsilon), \dots, (F_m, \varepsilon)$  to obtain a map  $G = (G_1, \dots, G_m)$  all of whose components obey the conclusions of Lemma 2.5. Let us write  $G_j(re^{i\theta}) = \sum_{n=-N(j)}^{N(j)} B_n(r)e^{in\theta}$ ,  $j = 1, 2, \dots, m$ . We apply Lemma 2.1 by defining

$$b_n(r) := B_{jn}(r), \quad j \neq 0, \quad b_0(r) := B_{j0}(r) + \eta,$$

for each  $j = 1, \dots, m$ , and obtain

$$\mathcal{A}_r^{(\eta)} := (\mathcal{A}_{1,r}^{(\eta)}, \dots, \mathcal{A}_{m,r}^{(\eta)}) : \overline{\text{Ann}}(0; r, 1) \rightarrow D^m,$$

$$\mathcal{A}_r^{(\eta)} \in \mathcal{O}[\text{Ann}(0; r, 1)] \cap \mathcal{C}[\overline{\text{Ann}}(0; r, 1); D^m] \text{ for every } \eta \in D(0; \delta)^m \text{ and } \forall r \in (0; 1),$$

where  $\delta > 0$  is chosen to be so small that :

- $\sup_{\partial D} |H_j| + \delta < 1$  for  $j = 1, \dots, m$ ;
- $\delta + \sum_{n=-N}^N |B_{jn}(r)|/r^n < 1 \forall r \in (0; 1]$  and  $j = 1, \dots, m$ ; and
- $H(D) + \eta \subset \Omega_1 \forall \eta \in D(0; \delta)^m$ .

As before, there exists a  $D^m$ -valued function  $\psi := (\psi_1, \dots, \psi_m) \in \mathcal{O}(D) \cap \mathcal{C}(\overline{D}; D^m)$  such that for each  $\eta \in D(0; \delta)^m$

$$\lim_{r \rightarrow 0^+} \mathcal{A}_r^{(\eta)}(\zeta_0) = \psi(\zeta_0) + \eta \text{ for each fixed } \zeta_0 \in D \setminus \{0\}.$$

Defining

$$H^{(\eta)} := (H_1(\zeta) + \eta_1, \dots, H_m(\zeta) + \eta_m) \quad \forall \eta = (\eta_1, \dots, \eta_m) \in D(0; \delta)^m,$$

$$\mathfrak{K}^{(\eta)} := \Gamma(\psi + \eta) \cup \left[ \cup_{0 < r < 1} \Gamma(\mathcal{A}_r^{(\eta)}) \right] \setminus \Omega_1,$$

we see that properties (1)–(3) in the proof of Theorem 1.1 hold for  $\{\mathcal{A}_r^{(\eta)}\}_{r \in (0,1)}$  and  $\psi$  in our new context – the only difference being that the relevant functions are vector-valued, and  $\eta$  varies in a polydisc  $D(0; \delta)^m$ . Therefore,  $\{\Gamma(\mathcal{A}_r^{(\eta)})\}_{r \in (0,1)}$  is a continuous family of analytic annuli attached to  $\Gamma(H^{(\eta)}) \cup (\partial D \times D^m)$ , which accumulate onto  $\Gamma(\psi + \eta)$  as  $r \rightarrow 0^+$ . The analytic annuli  $\Gamma(\mathcal{A}_r^{(\eta)})$  with  $r \approx 1$  are contained in  $\Omega_1$ . As before, the Kontinuitätssatz tells us that

$$(3.5) \quad U(\mathbf{P}) := \bigcup_{\eta \in \mathbf{P}} \Gamma(\psi + \eta) \subset \pi^{\Omega_1}(\widetilde{\Omega}_1),$$

where  $\mathbf{P}$  is any polydisc contained in  $D(0; \delta)^m$ .

Arguing exactly as before, we can find a sufficiently small polydisc  $\mathbf{P}_* \subset D(0; \delta)^m$  centered at the origin, an appropriately chosen domain  $\omega_2$  such that  $\overline{\Omega_2} \subset \omega_2 \subset \Omega_1$ , and a mapping  $\tilde{j} : U(\mathbf{P}_*) \cup \omega_2$  (here,  $U(\mathbf{P}_*)$  is as defined in (3.5) above) such that

- $\tilde{j} \in \mathcal{O}(U(\mathbf{P}_*) \cup \omega_2)$ , and
- $\tilde{j}|_{\omega_2} \equiv j|_{\omega_2}$ .

We use this mapping  $\tilde{j}$  exactly as in the previous two theorems to complete this proof.  $\square$

## 4. EXAMPLES

We begin by showing that one can construct a Hartogs-type configuration  $\Gamma(F) \cup (\partial D \times D)$  such that  $\sup_{\overline{D}} |F|$  is as large as we want and such that functions holomorphic in any small neighbourhood of this configuration extend to  $D \times D$  in the manner described in Theorem 1.1. A related *counterexample* to this sort of a phenomenon is the case when  $\Gamma(F)$  is a Wermer disc (see Example 4.2 below for a definition). In that case, *there is no analytic continuation*, provided  $\sup_{\overline{D}} |F|$  is sufficiently large. We explain under the heading Example 4.2 why this does not contradict Theorem 1.1

Theorem 1.3 is not true if  $F$  occurring therein is replaced by an arbitrary smooth,  $D^m$ -valued function. This is the content of Rosay's counterexample in [8]. Under Example 4.3 below, we discuss how Rosay's counterexample fails to meet the hypotheses of Theorem 1.3.

We begin with our first example.

**Example 4.1.** *An example showing that given any  $N \in \mathbb{N}$ , there is a  $F \in \mathcal{C}^\infty(\overline{D}; \mathbb{C})$  such that  $\sup_{\overline{D}} |F| > N$ ,  $\sup_{\partial D} |F| < 1$ , and such that small connected neighbourhoods of  $\Gamma(F) \cup (\partial D \times D)$  exhibit the analytic-continuation phenomenon described in Theorem 1.1.*

We are given  $N \in \mathbb{N}$ . Let  $\chi \in \mathcal{C}^\infty([0, \infty); [0, 1])$  be a smooth cut-off on  $[0, \infty)$  such that

$$\begin{aligned} \chi|_{[1/(N+1), \infty)} &\equiv \frac{N + (1/2)}{N + 1}, \\ \chi &\equiv 0 \text{ in a relatively open neighbourhood of } 0. \end{aligned}$$

We define

$$F(re^{i\theta}) := \frac{\chi(r)}{r} e^{-i\theta}.$$

Clearly

$$\begin{aligned} |F(e^{i\theta}/(N+1))| &= N + (1/2) > N, \\ |F(\zeta)| &= \frac{N + (1/2)}{N + 1} < 1 \quad \forall \zeta \in \partial D, \\ r|A_{-1}(r)| &= \chi(r) \leq \frac{N + (1/2)}{N + 1} < 1 \quad \forall r \in (0, 1]. \end{aligned}$$

These conditions imply that for any connected neighbourhood  $\Omega_1 \supset \Gamma(F) \cup (\partial D \times D)$ , any connected open set  $\Omega_2$  satisfying  $(\partial D \times D) \subset \overline{\Omega_2} \subset \Omega_1 \cap (\{|z| \geq 1\} \times D)$ , and for any  $f \in \mathcal{O}(\Omega_1)$ ,  $f|_{\Omega_2}$  extends holomorphically to  $D \times D$ .

Wermer presents an example of a function  $g \in \mathcal{C}^\infty(\overline{D})$  [6] with the property that  $\Gamma(g)$  is totally real, but  $g|_{\partial D} \equiv 0$ . This allows us to define a function  $F = Mg$  – where  $M > 0$  is sufficiently large – such that the configuration  $\Gamma(F) \cup (\partial D \times D)$  resists analytic continuation of the type described in Theorem 1.1. We now explain how the relevant  $F$  fails to satisfy the hypothesis of Theorem 1.1.

**Example 4.2.** *Wermer's disc.*

The graph of the function

$$g(z, \bar{z}) = \bar{z}(1 - |z|^4) + iz(1 - |z|^2)$$

is a totally-real surface in  $\mathbb{C}^2$ . This follows from an easy computation; see details in [6, Example 6.1]. Therefore, the domain

$$\mathcal{D}_\delta := \{(z, w) \in \mathbb{C}^2 : |z| < 1 + \delta, |w - g(z)| < \delta\}$$

is a pseudoconvex domain for all  $\delta > 0$  sufficiently small. Notice that  $\mathcal{D}_\delta \supset \partial D \times D(0, \delta)$ . Let  $\delta^* > 0$  be so small that  $D \times \{0\} \not\subseteq \mathcal{D}_{\delta^*}$  and  $\mathcal{D}_{\delta^*}$  is pseudoconvex. Then, for each domain  $\Omega_1 \supset \Gamma(g) \cup (\partial D \times D(0; \delta^*))$  such that  $\Omega_1 \subset \mathcal{D}_{\delta^*}$ , there would exist a function  $f \in \mathcal{O}(\Omega_1)$  such that  $f$  does *not* extend holomorphically to the bidisc  $D \times D(0; \delta^*)$ , because  $\mathcal{D}_{\delta^*}$  is a domain of holomorphy but does not contain  $D \times \{0\}$ . We now define

$$F(z) := \frac{1}{\delta^*} g(z), \quad \widetilde{\mathcal{D}}_{\delta^*} := \{(z, w) \mid (z, \delta^* w) \in \mathcal{D}_{\delta^*}\}.$$

By construction,  $\widetilde{\mathcal{D}}_{\delta^*}$  is a pseudoconvex domain that contains  $\Gamma(F) \cup (\partial D \times D)$  but does not contain  $D \times \{0\}$ . By our preceding remarks, the Hartogs-Chirka type configuration just constructed does not admit analytic continuation in the manner described in Theorem 1.1

Notice that  $F(re^{i\theta}) = A_{-1}(r)e^{-i\theta}$ , where

$$A_{-1}(r) = \frac{r}{\delta^*} \{ (1 - r^4) + i(1 - r^2) \}.$$

We will now show that

$$r|A_{-1}(r)| = \frac{r^2}{\delta^*} \{(1 - r^4)^2 + (1 - r^2)^2\}^{1/2} \gg 1 \quad \text{for some } r \in (0, 1],$$

whence Theorem 1.1 is inapplicable to the above configuration. For this purpose, we will need an upper bound for the quantity  $\delta^*$  introduced above, and we make the following

**Claim :**  $\delta^* < 0.0061$ . To see this, we refer to the Berndtsson-Słodkowski inequality – see [1, Prop.2.3/(b)] – determining when a surface of the form

$$\mathcal{S} = \{(z, w) \in \Omega \times \mathbb{C} \mid |w - G(z)| = e^{-u(z)}\},$$

(here  $\Omega$  is a domain in  $\mathbb{C}$ ,  $G$  and  $u$  are smooth functions, and  $u$  is real-valued) is pseudoconvex. The desired inequality is

$$(4.1) \quad \mathcal{S} \text{ is pseudoconvex} \iff -u_{z\bar{z}} \leq e^{2u}|G_{\bar{z}}|^2 - e^u|G_{z\bar{z}} + 2u_z G_{\bar{z}}|.$$

For  $\mathcal{D}_{\delta^*}$  to be pseudoconvex, we require that the surface  $\mathcal{S}_{\delta^*} := \{(z, w) \in D \times \mathbb{C} \mid |w - g(z)| = \delta^*\}$  be pseudoconvex. Applying (4.1) to the surface  $\mathcal{S}_{\delta^*}$  yields the following restriction on  $\delta^*$ .

$$0 < \delta^* \leq \frac{(1 - 3|z|^4)^2 + (1 - 2|z|^2)^2}{|z|\sqrt{36|z|^4 + 4}} \quad \forall |z| \leq 1.$$

In other words,

$$0 < \delta^* \leq \min_{r \in [0, 1]} \frac{(1 - 3r^4)^2 + (1 - 2r^2)^2}{r\sqrt{36r^4 + 4}},$$

and one can use any computational software package to show that the right-hand side of the above inequality is greater than 0.0061. Hence the claim.

One can also compute that  $\max_{r \in [0,1]} r^2 \{(1-r^4)^2 + (1-r^2)^2\}^{1/2} \approx 0.456$ . Thus

$$\max_{r \in [0,1]} r |A_{-1}(r)| > \frac{0.456}{0.0061} \gg 1,$$

which violates condition (1.1).

**Example 4.3.** *Rosay's counterexample.*

Rosay shows that one can find an arbitrarily small, strictly pseudoconvex neighbourhood  $\Omega$  of  $\partial D \times D^2$  and a  $D^2$ -valued function  $F$  such that  $\Gamma(F) \cup (\partial D \times D^2)$  is holomorphically convex. Specifically

$$\Omega := \{(z_1, z_2, z_3) \in \mathbb{C}^3 : \{(|z_1|^2 - 1)^2 + s_1 |z_2|^2\}^N + \left|\frac{z_3}{N}\right|^{2N} + \alpha \{(|z_1|^2 - 1)^2 + s_1 |z_2|^2 + |z_3|^2\} < s^{2N} + \alpha s^2\},$$

where

- $s > 0$  is small, and  $s_1 = s(1 - \delta) < s$  for a fixed, small  $\delta > 0$ , and
- One first chooses  $N$  large enough that  $s_1^{2N} + 1/N^{2N} < s^{2N}$ , and then chooses  $\alpha$  sufficiently small so as to ensure that  $\partial D \times D^2 \Subset \Omega$ .

Write  $F = (F_1, F_2)$ . In Rosay's construction

$$F_1(re^{i\theta}) := \kappa \chi(r) e^{i\theta},$$

where  $\chi \in C^\infty[0, 1]$  with  $0 \leq \chi \leq 1$ , such that  $\chi \equiv 1$  off a small relative neighbourhood of  $0 \in [0, 1]$  and  $\chi \equiv 0$  on a smaller neighbourhood of  $0$ . The quantity  $\kappa$  will be described presently. The function  $F_2$  is required to be identically zero in an open set contained in  $\{re^{i\theta} : r \in \text{supp}(\chi)\}$ , and to satisfy  $\partial F_1 / \partial \bar{z}_1 \neq 0$  wherever  $F_1 \equiv 0$ . Therefore,  $F_2$  will have negative Fourier modes.

Our interest is in examining  $F_1$ . The constant  $\kappa$  is so chosen that

$$\Gamma(F) \cap \bar{\Omega} = \{(Re^{i\theta}, \kappa e^{i\theta}, 0) : \theta \in [0, 2\pi)\},$$

and such that  $\Gamma(F) \cap \bar{\Omega}$  is a complex-tangential curve in the surface  $\partial\Omega \cap \{z_3 = 0\}$ . Write  $z_j := x_j + iy_j$ ,  $j = 1, 2$ . It is easy to determine what the magnitudes of  $\kappa$  and  $R$  (which is close to 1) should be by visualizing  $\omega := \bar{\Omega} \cap (\mathbb{R}^2 \times \{0\})$ . Then  $(R, \kappa)$  are the coordinates of the point of tangency, in the first quadrant, of the line through the origin that is tangential to  $\partial\omega$  (then, the complex span of this line contains the tangent line to the curve  $\Gamma(F) \cap \bar{\Omega}$ ). By construction, the point  $(1, s/s_1) \in \partial\omega$  lies *below* the line just described, whence the line  $x_1 = x_2$  lies below this line, in the first quadrant. Thus, in the notation of Theorem 1.3

$$\kappa/R := A_{11}(R)/R > 1,$$

whence, by construction

$$A_{11}(r)/r > 1 \quad \forall r \in \chi^{-1}\{1\}.$$

This violates the condition (1.2).



## REFERENCES

- [1] B. Berndtsson, *Levi-flat surfaces with circular sections*, Several Complex Variables, Math. Notes (John Erik Fornaess ed.), Princeton University Press, Princeton, NJ (1993), pp. 136-159.
- [2] G. Bharali, *Some generalizations of Chirka's extension theorem*, Proc. Amer. Math. Soc. **129** (2001), 3665-3669.
- [3] E.M. Chirka, *Generalized Hartogs' lemma and non-linear  $\bar{\partial}$ -equation*, Complex Analysis in Contemporary Mathematics (E.M. Chirka, ed.), FAZIS, Moscow (in Russian) (2002).
- [4] E.M. Chirka and J.-P. Rosay, *Remarks on the proof of a generalized Hartogs lemma*, Ann. Pol. Math. **70** (1998), 43-47.
- [5] E.M. Chirka and E.L. Stout, *A Kontinuitätssatz*, Topics in Complex Analysis (P. J ac obczak, W. Ple niak, eds.), Banach Center Publications, Warsaw, **31** (1995), pp. 143-150.
- [6] L. H ormander and J. Wermer, *Uniform approximation on compact sets in  $\mathbb{C}^n$* , Math. Scand. **23** (1968), 5-21.
- [7] S.M. Ivashkovich and V.V. Shevchishin, *Deformations of noncompact complex curves, and envelopes of meromorphy of spheres* (in Russian), Mat. Sb. **189** (1998), 23-60.
- [8] J.-P. Rosay, *A counterexample related to Hartogs' phenomenon (A question by E. Chirka)*, Michigan Math. J. **45** (1998), 529-535.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 525 EAST UNIVERSITY ANN ARBOR, MI 48109  
E-mail address: [barrett@umich.edu](mailto:barrett@umich.edu)

E-mail address: [bharali@umich.edu](mailto:bharali@umich.edu)